

# HOW TO OBTAIN YOUR MEDICAL MARIJUANA CARD

## STEP 1

### REGISTER

- Visit the Patient and Caregiver Registry: [medicalmarijuana.pa.gov](https://medicalmarijuana.pa.gov) and create a patient profile.
- Ensure that the information you enter matches your state issued ID EXACTLY.
- Type in all CAPS.
- Do not use spaces or dashes between identification numbers.
- Do not use any punctuation in your address.

## STEP 2

### GET CERTIFIED BY A PHYSICIAN


- Visit a physician that is registered with the PA medical marijuana program.
- Receive a certification that you suffer from one of the conditions that qualifies you to receive a medical marijuana ID card.

## STEP 3

### MAKE PAYMENT

- Return to the Patient and Caregiver Registry and pay the \$50 fee for your medical marijuana ID card.
- Wait for your ID card to arrive before obtaining medical marijuana from an approved PA dispensary.





## HOW TO OBTAIN YOUR MEDICAL MARIJUANA CARD



### CAREGIVER INSTRUCTIONS

- STEP 1:** Visit the Patients and Caregiver Registry and create a caregiver profile. Do not use any punctuation in your address.
- STEP 2:** Complete a criminal background check.
- STEP 3:** Complete an 'authorization to designate a third-party caregiver' form.
- STEP 4:** Once the patient has their physician certification, return to the Patient and Caregiver Registry, and pay the \$50 fee for your medical marijuana ID card.

## QUALIFYING CONDITIONS

- Amyotrophic lateral sclerosis;
- Autism;
- Cancer, including remission therapy;
- Crohn's disease;
- Damage to the nervous tissue of the central nervous system (brain-spinal cord) with objective neurological indication of intractable spasticity, and other associated neuropathies;
- Dyskinetic and spastic movement disorders;
- Epilepsy;
- Glaucoma;
- HIV / AIDS;
- Huntington's disease;
- Inflammatory bowel disease;
- Intractable seizures;
- Multiple sclerosis;
- Neurodegenerative diseases;
- Neuropathies;
- Opioid use disorder for which conventional therapeutic interventions are contraindicated or ineffective, or for which adjunctive therapy is indicated in combination with primary therapeutic interventions;
- Parkinson's disease;
- Post-traumatic stress disorder;
- Severe chronic or intractable pain of neuropathic origin or severe chronic or intractable pain;
- Sickle cell anemia; and
- Terminal illness.